

HIV seroprevalence and seroincidence among drug users in Buenos Aires, Argentina.

Vila, Marcelo, Pando, María de los Ángeles, Ralón, Gonzalo, Patracca, Claudia, Vignoles, Moira, Ávila, María Mercedes, Salomón, Horacio y Rossi, Diana.

Cita:

Vila, Marcelo, Pando, María de los Ángeles, Ralón, Gonzalo, Patracca, Claudia, Vignoles, Moira, Ávila, María Mercedes, Salomón, Horacio y Rossi, Diana (Diciembre, 2008). *HIV seroprevalence and seroincidence among drug users in Buenos Aires, Argentina. XVII International AIDS Conference. International AIDS Society, Ciudad de México.*

Dirección estable: <https://www.aacademica.org/gonzalo.ralon/27>

ARK: <https://n2t.net/ark:/13683/pcqZ/0YM>



Esta obra está bajo una licencia de Creative Commons.
Para ver una copia de esta licencia, visite
<https://creativecommons.org/licenses/by-nc-nd/4.0/deed.es>.

Acta Académica es un proyecto académico sin fines de lucro enmarcado en la iniciativa de acceso abierto. Acta Académica fue creado para facilitar a investigadores de todo el mundo el compartir su producción académica. Para crear un perfil gratuitamente o acceder a otros trabajos visite:
<https://www.aacademica.org>.



XVII INTERNATIONAL AIDS CONFERENCE

3-8 August 2008 | Mexico City

CD-ROM Publication

Track C - HIV seroepidemiologic studies

CDC0140 - HIV seroprevalence and seroincidence among drug users in Buenos Aires, Argentina

M.E. Vila¹, M. Pando², G. Ralón¹, C. Petracca³, M. Vignoles², M. Avila², H. Salomón², D. Rossi¹

¹Intercambios Asociacion Civil. Fogarty International Training and Research project D43 TW001037-06 (Mount Sinai / New York State / Argentina HIV Prevention), Buenos Aires, Argentina, ²Centro Nacional de Referencia para el Sida. Facultad de Medicina, Universidad de Buenos Aires, Buenos Aires, Argentina, ³Programa Municipal de Sida de San Martín. Provincia de Buenos Aires, Buenos Aires, Argentina

Background: Argentina has a high proportion (32%) of AIDS cases among injecting drug users (IDUs). Few studies show incidence and prevalence in IDUs and in non-injecting drug users (NIDUs). This information is necessary to understand the trend of the epidemic and to design adequate responses. Part of the study was funded by The Global Fund (Sub-Project N°112).

Estimate HIV seroprevalence and seroincidence among current drug users (IDUs and NIDUs) in the Buenos Aires Metropolitan Area and to understand the association with their sexual and drug use practices.

Methods: Between Oct2006/Oct2007, 358 current cocaine/amphetamine users were recruited; 53% were recruited in drug treatment centers and 47% through harm reduction programs in the neighborhoods where they live. Informed consent, a structured questionnaire and HIV diagnosis and counseling were carried out. Seroincidence was calculated through the detune assay.

Results: Sample was 62% male; mean age: 30 years old. HIV Prevalence: 6.7% (95% CI 4.4-9.6) without difference by sex. In a bivariate analyses, variables associated with high level of prevalent infection (OR>1) were: IDUs in last year: 6.5 (1.9-22.5); anal sex between men in last year: 5.4 (2.0-14.4); over 25 years old: 7.6 (1.8-33.0) HIV seroincidence was 3.2 per 100 person-years at risk (95% CI 1.7-5.7) without difference by sex.

All recent infections were among non-injecting drug users, 50% among men with history of MSM and 50% among women.

Conclusions: Variables associated with high rate of prevalent infection were previous experience with injection drug use, unprotected anal sex among men, and greater age. It is necessary to understand more about sexual and drug use practices among non-injecting drug users in order to develop adequate prevention strategies.

Presenting author email: mvila@intercambios.org.ar